

Schizophrenia



Schizophrenia is marked by positive symptoms

Schizophrenia is one of Lundbeck's focus disease areas, and is a chronic, severe and disabling psychiatric disorder. The disease is characterized by distortions in thinking, perception, emotions, language, sense of self and behaviour. Common experiences include hearing voices and delusions. Schizophrenia affects both men and women, although men tend to develop the condition slightly earlier in life.

hallucinations delusions



blunted emotions social withdrawal

The World Health Organization estimates that over

21
million people

suffer from schizophrenia, making it one of the top 10 causes of disability worldwide.¹



Schizophrenia

is among the most financially costly illnesses in the world ^{3, 4}

Schizophrenia affects people regardless of race, culture or social class. It typically starts in early adulthood around age

20

but it can develop at any age.2

The risk of an individual developing schizophrenia during his or her lifetime is approximately



- World Health Organization. Schizophrenia fact sheet. 2016.
 (http://www.who.int/mediacentre/factsheets/fs397/en/)
- Ochoa S, Usall J, Cobo J, Labad X, Kulkarni J. Gender differences in schizophrenia and first-episode psychosis: a comprehensive literature review. Schizophr Res Treatment 2012; 2012: 916198.
- Lindström E, Eberhard J, Neovius M, Levander S. Costs of schizophrenia during 5 years. Acta Psychiatr Scand Suppl 2007;116 (435):3340.
- Rössler W, Salize HJ, van Os J, Riecher-Rössler A. Size of burden of schizophrenia and psychotic disorders. Eur Neuropsychopharmacol 2005; 15 (4): 399409.
- Tsuang MT, Farone SV. Schizophrenia. Second edition. Oxford University Press Inc., New York: 2005.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders, 5th edition (DSM-5). Washington, D.C.: American Psychiatric Association; 2013.